

Children's Cough & Cold Liquid

Drug Facts

Active ingredients *Purpose* (in each 5 mL)

Diphenhydramine HCl 12.5 mg.....	Antihistamine/ Cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses

- temporarily controls cough due to minor throat and bronchial irritation and relieves nasal congestion as may occur with a cold
- temporarily relieves the following symptoms due to hay fever or other upper respiratory allergies:
 - sneezing
 - runny nose
 - itchy, watery eyes
 - itchy nose or throat

Warnings

Do not use

- to make a child sleepy
- with any other product containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if

- you are taking sedatives or tranquilizers.

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with fever, rash or persistent headache. These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter

Age (year)	Dose (mL)
adults and children 12 years of age and over	10 mL every 4 hours
children 6 to under 12 years of age	5 mL every 4 hours
children 4 to under 6 years of age	do not use unless directed by a doctor
children under 4 years of age	do not use

Other information

- each 5 mL contains: sodium 10 mg
- tamper evident: do not use if neckband on bottle cap is broken or missing
- dosing cup provided
- store between 15-30°C (59-86°F)
- Keep carton for full directions for use

Inactive ingredients anhydrous citric acid, anhydrous trisodium citrate, carboxymethylcellulose sodium, edetate disodium, FD&C Blue #1, FD&C Red #40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

Questions? 1-888-963-3382

You may also report side effects to this phone number.